

# New type of extreme value statistics

D.B. Saakian

Yerevan Physics Institute, Alikhanian Brothers St. 2, Yerevan 375036,  
Armenia.

## Abstract

We investigate the extreme value statistics connected with the dilute Random Energy Model with integer couplings. New universality class is found.

Extreme value statistics [1] statistics play an important role in statistical mechanics and theory of probability [2-5]. They describe the minimal and maximal bounds for the distribution of a large number of independent random quantities, and it is the main reason of their recent applications in statistical physics of disordered systems.

Usually one considers sequence of  $M = 2^N$  independent random variables  $E_i$  with the same distribution

$$\rho(E) = \langle \delta(E_i - E) \rangle \quad (1)$$

Let us find minimal and maximal members  $E_{min}$  and  $E_{max}$  of our sequence, then consider their distribution at the limit of large N. When  $\rho(E)$  is independent from N, at  $N \rightarrow \infty$  limit only 3 classes of universalities has been found [1] for the distribution of  $\rho(E_{min})$ . After rescaling of  $x \equiv E_{min}$  one has for them:

$$P_1(x) = 1, x \geq 0, P_1(x) = 1 - \exp[-(-x)^k] \quad x \leq 0 \quad (2)$$

$$P_2(x) = 0, x \leq 0, P_2(x) = 1 - \exp[-x^k] \quad x \geq 0 \quad (3)$$

$$P_3(x) = 1 - \exp[-e^x] \quad (4)$$

Now we are going to consider another (physically interesting) case, when our  $\rho(E)$  is a superposition of N random variables.

In the present paper we will investigate extreme value statistics which are generated by the dilute-coupling REM. Surprisingly, we got new types of them due to physical peculiarities of the model.

The dilute coupling REM is described by the hamiltonian

$$H = - \sum_{(i_1 \dots i_p)=1}^{\alpha N} j_{i_1 \dots i_p} s_{i_1} \dots s_{i_p} \quad (5)$$

where only  $\alpha N$  quenched bonds are nonzero, and they take values  $\pm 1$  with equal probability.  $s_i = \pm 1$  are the Ising spins.

In the limit of large p there is a factorization in probability of different energy

levels and we have  $2^N$  independent random variables (energies) with the same probability distribution  $\rho(E)$ . For our choice of integer couplings we have for distribution  $\rho(E)$ .

$$\rho(E) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} dk e^{N\varphi(k) - kE} \quad (6)$$

where

$$\varphi(k) = \alpha \ln \cosh(k) \quad (7)$$

This representation is correct for

$$|E| < \alpha \quad (8)$$

Otherwise  $\rho(E)$  is strictly zero.

For the case  $\alpha < 1$  we should keep in mind, that there are  $2^{\alpha N}$  different classes of energy configurations with

$$2^{(1-\alpha)N} \quad (9)$$

equal energies in each one. Energies from different classes are distributed independently according to (6).

At low temperatures the system is frozen in minimal energy configuration (chosen among the all  $2^N$  ones). That's why the physics of REM is connected closely with extreme value statistics problem.

As known, several phases are possible in the considered model. For  $\alpha > 1$  there is a phase transition to the spin glass phase and at high temperatures the system is in the paramagnetic phase. For the case  $\alpha < 1$  only the paramagnetic phase is possible. The limiting case  $\alpha \sim 1$  corresponds to the border between 2 phases.

We hope, that different universality classes are possible, connected with those phases. Let us consider  $2^N$  independent random variables with (6), where  $\varphi(k)$  is even analytical function with respect to  $k$ . It is just enough for the most of situations. For the case of hamiltonian (5) one has, that

$$\lim_{k \rightarrow \infty} \varphi(k) = \alpha(k - \ln 2) + o(\exp(-2k)) \quad (10)$$

We are considering only distributions, which have this asymptotic form, or when corrections are polynomial

$$\lim_{k \rightarrow \infty} \varphi(k) = \alpha(k - \ln 2) + o(k^{-\beta}) \quad (11)$$

We are going to consider now the extreme value statistics for  $2^N$  random variables with our representation for  $\rho(k)$ .

Let us define the probability  $\theta(E)$ , as the integral of these expressions in the interval  $[-\infty, E]$ . We have

$$\theta(E) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} dk e^{N\varphi(k) - kE} / k \quad (12)$$

Here our integration loop overpasses the point 0 from the left size.

Let us find the distribution of the minimal bound. We take one variable  $E_i$  in our interval  $[-\infty, E]$ , the others to right. As the first one could be chosen from the all  $2^N$ , we have an expression [4]

$$P(E) = 2^N \rho(E) (1 - \theta(E))^{2^N - 1} = -\frac{d}{dE} (1 - \theta(E))^{2^N} \quad (13)$$

We need to study the expression  $\theta(E)$  in the region, when it is small. Here

$$(1 - \theta(E))^{2^N} \approx \exp[-2^N \theta(E)] \quad (14)$$

Its density is nonzero only near the  $E = E_c$ . We need to find asymptotic of function  $\theta(E)$  near the  $E_c$ , defined by condition

$$2^N \theta(E_c) = 1 \quad (15)$$

For the different choices of  $\alpha$  we have to consider different asymptotic regims for  $\theta(E)$ . In this way different universality classes arose.

While calculating (12) for the saddle point we have an equation

$$\varphi'(k) = -\frac{E}{N} \quad (16)$$

There are three possibilities for  $E_c$ . The first one

$$|E_c| < \alpha N \quad (17)$$

has the same statistics, as in the case when  $\varphi(k)$  goes to infinity faster than linearly (this is the case of ordinary REM [4] with normal distribution of energies). Two others are possible also:

$$|E_c| \sim \alpha N \quad (18)$$

$$|E_c| > \alpha N \quad (19)$$

In the last case we should take complex solutions of (16). Let us first calculate the first case (18). Two expressions for the function  $\theta(E)$  coincide (at the limit of large N) and we have

$$\theta(E) = \frac{1}{k \sqrt{2\pi N \varphi''(k)}} e^{N \varphi(k) - kE} \quad (20)$$

One can write the expression for the desired probability distribution for the minimum of our  $2^N$  random variables:

$$e^{\left(-\frac{2^N}{k \sqrt{2\pi N \varphi''(k)}} e^{N \varphi(k) - kE}\right)} \frac{d}{dE} \frac{1}{k \sqrt{2\pi N \varphi''(k)}} e^{N \varphi(k) - kE} \quad (21)$$

For our asymptotic form (15) transforms into

$$\frac{2^N}{k\sqrt{2\pi N\varphi''(k)}}e^{N\varphi(k)-kE_c} = 0 \quad (22)$$

where  $k$  is defined from (16). For the  $E < E_c$  our probability is almost one, and above it is almost zero. For a small deviation

$$u \equiv (E - E_c)k_c \ll N \quad (23)$$

we obtain Gumbel distribution

$$P(u) = \exp(u - \exp(u)), \quad (24)$$

where  $u$  changes in the interval  $-\infty, \infty$ . Let us now consider the statistics for limiting situation, when  $\alpha \sim 1$ . We shall consider the distribution (7), but the corresponding limiting statistics is the same for the whole class of exponential corrections. At the saddle point

$$\tanh(k) = E/(\alpha N) \quad (25)$$

we have

$$\begin{aligned} \theta(E) &= \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{\alpha N \ln \cosh(k) - kN} / k \approx \\ &= \frac{\cosh(k)}{k\sqrt{N2\pi}} e^{-\alpha N g(E/(\alpha N))} \end{aligned} \quad (26)$$

where

$$g(x) = \frac{1+x}{2} \ln \frac{1+x}{2} + \frac{1-x}{2} \ln \frac{1-x}{2} \quad (27)$$

Let us do notation

$$N_1 = N\alpha, \frac{E}{\alpha N} = 1 - \epsilon \quad (28)$$

We are considering the case, where  $\frac{N_1}{N} - 1 \equiv \epsilon_1 \rightarrow 0$ . In the limit  $\epsilon \rightarrow 0$  we have

$$g(1 - \epsilon) \rightarrow \ln 2 - \frac{\epsilon}{2} \ln\left(\frac{2}{\epsilon}\right), \frac{1}{\cosh(k)^2} \sim \epsilon \quad (29)$$

$$\theta(E) = \frac{1}{\ln(\frac{2}{\epsilon})\sqrt{N\pi\epsilon}} \exp\left(-N_1 \ln 2 + \frac{N_1\epsilon}{2} \ln(2/\epsilon)\right) \quad (30)$$

Let us check, when such a behaviour is correct. We should take condition

$$\frac{N}{\cosh^2(k)} \sim N\epsilon \gg 1 \quad (31)$$

Now let us consider the region  $N\epsilon \sim 1$ . Of course, at the saddle point we cannot approximate our integral by the gaussian one for our values of  $\epsilon$ , since the second derivative tends to zero. We should directly calculate integral.

For the complex  $k \equiv k_1 + ik_2$  we have

$$\begin{aligned} \ln \cosh(k) &= \cosh k_1 + 1/2 \ln[1 - (1 - \tanh^2 k_1) \sin^2 k_2] \\ &+ i \arctan[\tan k_2 - (1 - \tanh k_1) \tan k_2] \end{aligned} \quad (32)$$

This is an exact expression. Let us consider the case  $k_1 \rightarrow \infty$ . In this limit

$$\ln \cosh(k) = k_1 - \ln 2 + e^{-2k_1}, \quad \tanh k_1 = 1 - 2e^{-2k_1} \quad (33)$$

So we have to choose the position of  $k_1$ , while performing integration

$$\frac{1}{2\pi i} \int_{k_1 - i\infty}^{k_1 + i\infty} e^{\alpha N [\ln \cosh(k) - (1-\epsilon)k]} / k \quad (34)$$

The saddle point equation gives the choice

$$k_1 = \frac{1}{2} \ln\left(\frac{2}{\epsilon}\right) \quad (35)$$

With such a choice of  $k_1$  we get

$$\begin{aligned} N_1 \ln \cosh(k) - N_1(k_1 + ik_2) + N_1\epsilon(k_1 + ik_2) \\ \approx N(\epsilon/2 + \epsilon/2 \ln 2N/\epsilon - \epsilon \sin k_2 + i\epsilon(k_2 - \sin k_2 \cos k_2)) \end{aligned} \quad (36)$$

Accuracy of this expression is  $\exp(-2k_1) \sim \epsilon$ . Our expressions are valid in a region

$$N \gg \epsilon \quad (37)$$

We get

$$\begin{aligned} \theta(E) &\approx \exp\left(-N_1 \ln 2 + \frac{N\epsilon}{2} \ln\left(\frac{2}{\epsilon}\right)\right) \\ &\frac{1}{\pi} \int_0^\infty dk_2 (k_1^2 + k_2^2)^{-1} e^{-\epsilon \sin k_2^2 k_1} \cos(\epsilon(k_2 \\ &- \sin k_2 \cos k_2)) + k_2 \sin(\epsilon(k_2 - \sin k_2 \cos k_2)) \end{aligned} \quad (38)$$

For the limit of small  $\epsilon$  last expression transforms into

$$\theta(E) = \exp\left(-N_1 \ln 2 + \frac{N\epsilon}{2} \ln(2/\epsilon)\right) \quad (39)$$

We have for the desired distribution for  $u = (\frac{E}{\alpha N} - 1)N$

$$P(u) = \left(-1 + \ln \frac{N}{u}\right) \exp\left[u \ln \frac{N}{u} - \exp\left(u \ln \frac{N}{u}\right)\right] \quad (40)$$

Such distribution is correct in the interval  $-\infty < u < 0$ . What about negative values of  $u$ ? For the choice (7) with integer  $\alpha N$  we have, that there is a strict bound for energy distribution equal to  $N\alpha$ , so for that case we can put zero probability for negative values of  $u$ . This expression should exist for any bounded discrete distribution of energies.

In [5] authors consider fully connected version of REM, here perhaps could be only classic distribution among the (2)-(4). The case of bounded continuous distribution brings to (2)-(3).

This work was supported by Fundacion Andes grant c-13413/1. I would like to thank E. Vogel for hospitality in UFRO.

## References

- [1] E. J. Gumbel, *Statistics of Extremes*, Columbia University Press, (1958);  
J. Galambos, *The asymptotic Theory of Extreme Order Statistics*, R.E. Krieger Publishing Co., Malabar, Florida (1987).
- [2] J.P. Bouchaud, A. Georges, Phys. Rep. **195** 127 (1990).
- [3] M. Marsili J. Phys. **29A** (1996) 5405.
- [4] J.P. Bouchaud, M. Mezard, J. Phys. **30A** (1997) 7997.
- [5] B. Derrida, Phys. Rev. Lett. **45** (1980) 79.
- [6] D. Dominicus, P. Mottishaw, J. Phys., **A20**(1987)L267
- [7] A.E. Allahverdyan, D.B. Saakian, Nucl. Phys **v.B498** (1997)604.
- [8] D.J. Gross, M. Mezard, Nucl. Phys. **B240** 431 (1984).